

Anastomosis of Free Flap Pedicle to Great Vessels

Jalaluddin Khoshnevis^{1*}, Terifeh Dashti², Mohammad Ebrahimi¹, Eznollah Azargashb³,
Mohamadreza Kalantar Motamedi¹

1. General and Vascular Surgery, Shohada-Tajrish Medical Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran
2. Health Service Management, Clinical Research Development Center of Shohada-Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran
3. Department of Community Medicine, Shahid Beheshti University of Medical Sciences and Health Services, Shohada-Tajrish Hospital, Tehran, Iran

ABSTRACT

BACKGROUND

Free Flaps are viable option to cover the tissue defect. Pedicle anastomosis to vessel branches has excellent result. In some situations which there is a possibility of flap failure like shortage of vessel branches, possibility of pedicle kinking or need to vein graft, anastomosis to great vessels is justified.

METHODS

Six patients were allocated to study. Five cases for free jejunal flap and one case for free latissimus flap. In free jejunal flap group, pedicle anastomosis was performed as an end-side fashion to common carotid artery and internal jugular vein and in free latissimus flap, pedicle was anastomosed as an end-side fashion to superficial femoral artery and superficial femoral vein. Follow up was regular up to 20 years.

RESULTS

In free jejunal flap group, there were three female and two male with age from 30 to 59 years. The sixth case was a thirteen years old male with flexion contracture of right knee who underwent free latissimus flap. Follow up was regular for 20 years. All flaps survived, and good functional result was obtained in all except one.

CONCLUSION

Choosing great vessels as one side of anastomosis is safe and can be done as a primary approach due to technical demand or as a final resort when there is shortage of side branches.

KEYWORDS

Free jejunal, Flap, Great vessels, Latissimus

Please cite this paper as:

Khoshnevis J, Dashti T, Ebrahimi M, Azargashb E, Kalantar Motamedi MR. Anastomosis of Free Flap Pedicle to Great Vessels. *World J Plast Surg* 2018;7(3):351-356. doi: 10.29252/wjps.7.3.351.

INTRODUCTION

Free jejunal flap was the first free flap described in the literature. From Seidenberg report at 1959 to date, remarkable advances have been achieved in the technique with overall success of 95-97%.¹⁻³ The tube lengths of up to 30 cm has been harvested and replaced lost esophagus successfully.⁴ The second jejunal branch is usually the best site for flap harvesting providing arterial pedicles up to 3-4 mm in diameter. All arterial and venous anastomosis were done to side branches of subclavian or carotid artery and jugular veins.¹⁻⁴

*Corresponding Author:

Jalaluddin Khoshnevis, MD;
General and Vascular Surgery,
Shohada-Tajrish Medical Center,
Shahid Beheshti University of Medical
Sciences,
Tehran, Iran.

Tel/Fax: +98-21-22721144

Email: khoshnevis.j@gmail.com

Received: August 30, 2017

Revised: July 25, 2018

Accepted: August 2, 2018

Few anastomosis were done to external carotid artery.⁵ Vein grafts were accused as one of the causes of flap failure.⁶ Also failures was reported from venous kinking.⁶ Choosing the recipient vessels is still a challenge.⁷⁻¹⁰ Many factors influence the choice of recipient vessels. History of head and neck radiation, surgery, and type of reconstruction dictates the choice of recipient vessels. It goes without saying tributaries are the first choice avoiding the possible complication by choosing the great vessels as recipient vessels. Here, we wish to introduce safety of the anastomosis to great vessels as recipient vessels to expand the choices and avoid additional dissections in new or complicated cases.

MATERIAL AND METHODS

Six patients were allocated to free flap with anastomosis of its pedicle to great vessels. Five out of six underwent free jejunal flap. After harvesting of the flap, side-side anastomosis was undertaken between jejunum and pharyngoesophagus. Then, we attended the position of the pedicles, which at the most of them were lying on the common carotid artery

and internal jugular vein. Totally, 100 IU/KG of heparin was injected as IV bolus before harvesting of flap.

The pedicle opening was spatulated three times of their diameter and anastomosed as an end-side fashion to common carotid artery and internal jugular vein with prolene 8/0 under 3x loupe magnification. Little maneuver such as pedicle dissection at the mesentery or orientation of arteriotomy and venotomy may be required for good adjustment. Proximal opening of the jejunal tube was spatulated and anastomosed to the side of pharyngotomy or closed primarily. The distal ostomy was closed two month later after complete healing of the wound.

RESULTS

We had excellent results in four of five free jejunal flaps and complete relief of dysphagia. In one patient, the failure was due to severe scar tissue of pharyngeal region due to burn injury and breakdown of pharyngojejunal anastomosis and consequently the wound. Despite this problem, the flap survived for two months. The sixth case who underwent free latissimus flap for knee contracture had good results with normal walking (Table 1).

Table 1: Demographic characteristics of the patients.

Case	Sex	Age (year)	Pathology	Type of flap	Vessel anastomosis to	Follow up (year)	Complications	Figure
1	F	59	Anastomosis stricture after orringer operation.	Free jejunal flap	Common carotid artery internal jugular vein	5	No	1
2	F	55	Unresectable esophageal web through endoscopy	Free jejunal flap	Common carotid artery internal jugular vein	10	No	2 and 3
3	F	50	Stricture due to epidermolysis bullosa	Free jejunal flap	Common carotid artery internal jugular Vein	20	No	4 and 5
4	M	47	Stricture due to epidermolysis bullosa	Free jejunal Flap	Common carotid artery internal jugular vein	8	No	6 and 7
5	M	30	Stricture due to caustic agent	Free jejunal flap	Common carotid artery internal jugular vein	2 Months	Wound dehiscence	-
6	M	13	Knee contracture after burn injury	Free latissimus flap	Superficial femoral artery superficial femoral vein	10	No	8-10




Fig. 1: Sever stricture (arrow).




Fig. 2: Esophageal web (arrow).




Fig. 3: Proximal end (white arrow) ,distal end (black arrow).




Fig. 4: End-side pedicle vein to internal jugular vein anastomosis (white arrow) and end-side pedicle artery to common carotid artery anastomosis (black arrow).



Fig. 5: Nail destruction and web.




Fig. 6: Blind loop during barium swallow (white arrow) and during evacuation by reverse peristalsism (black arrow) after 8 years.

DISCUSSION

Free jejunal flaps were used as tubular flap to substitute pharyngoesophageal defect or were opened to replace the mouth floor defect. In our cases, the jejunum was used as a tube, which was opened by its side and anastomosed to the side

esophagotomy. After side-side esophagojejunal anastomosis, the pedicle lie was evaluated just the moment, which laid in the middle of common carotid artery and internal jugular vein. Many authors ignored this ideal situation for anastomosis due to potential complication of anastomosis to common carotid artery and




Fig. 7: Esophageal stricture (arrows).



Fig. 8: Flexion contracture of knee (arrow).




Fig. 9: Anastomosis to SFA (white arrow) and SFV (black arrow)



Fig. 10: Latissimus dorsi flap after vein anastomosis revision (arrow).

makes tributaries as first recipient vessels.¹⁰⁻¹⁴

This is a true logic but when the tributaries have been destroyed due to previous surgery and radiation, you may go far for seeking

new recipient vessel. This needs more times and dissections and many need vein graft to accomplish the procedure, which ultimately may lead to more failure.¹⁵⁻¹⁸ Our approach to

common carotid artery as first recipient showed to be a safe one and the results was comparable to other authors.¹⁹⁻²¹ We had no flap failure, but only a wound dehiscence. The flap survived for two months before the next plan. Side-side esophagojejunal anastomosis proved to be a best one, because we had no dysphagia and stenosis during their long follow-up.

Considering two cases of epidermolysis bullosa cases who were prone to stenosis adds to the merit of this type of anastomosis. End to distal side esophagotomy prohibited due to more dysphagia occurrence.²¹ In the sixth case who had the history of sever burn injury and four surgeries to treat the knee contracture, we decided to treat with free latissimus flap. There were paucity of recipient tributary and thick femoral vein wall due to inflammation. We had no choice except end-to-side anastomosis to femoral artery and vein.

After flap congestion, the wound was explored and a small thrombosis was removed behind the femoral vein anastomosis. Drainage problem relieved by punching the vein and re-anastomosis. Despite this problem, the flap survived and knee contracture was relieved. After ten years, walking was normal. At this site, also the vessel recipient of choice was the tributaries.^{22,23} Few studies were found to mention anastomosis to the major vessels as recipient.² All of them had good results.²⁴⁻²⁶ We look forward for further cases to find a new place for this type of anastomosis at this site in algorithm.

In our study, anastomosis of pedicle of free flaps to great vessels was justified. Though anastomosis to branches has its own merit, there is situation like paucity of branches due to burn, surgery, radiotherapy or need for a vein graft or possibility of kinking in which the incidence of flap failure rises. So direct anastomosis of the pedicle to the side of common carotid artery and internal jugular vein was shown to be safe and worthy.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

- 1 Seidenberg B, Rosenak SS, Hurwitt ES, Som ML. Immediate reconstruction of the cervical

- esophagus by a revascularized isolated jejunal segment. *Ann Surg* 1959;**149**:162-71.
- 2 Numajiri T, Sowa Y, Nishino K, Fujiwara H, Nakano H, Shimada T, Hisa Y. Does a vascular supercharge improve the clinical outcome for free jejunal transfer? *Microsurgery* 2013;**33**:169-72.
- 3 Walker RJ, Parmar S, Praveen P, Martin T, Pracy P, Jennings C, Simms M. Jejunal free flap for reconstruction of pharyngeal defects in patients with head and neck cancer-the Birmingham experience. *Br J Oral Maxillofac Surg* 2014;**52**:106-10.
- 4 Nagasao T, Shimizu Y, Kasai S, Hatano A, Ding W, Jiang H, Kishi K, Imanishi N. Extension of the jejunum in the reconstruction of cervical oesophagus with free jejunum transfer using the thoracoacromial vessels as recipients. *J Plast Reconstr Aesthet Surg* 2012;**65**:156-62.
- 5 Okazaki M1, Asato H, Sarukawa S, Takushima A, Nakatsuka T, Harii K. Availability of end-to-side arterial anastomosis to the external carotid artery using short-thread double-needle micro suture in free-flap transfer for head and neck reconstruction. *Ann Plast Surg* 2006;**56**:171-5.
- 6 Kroll SS, Schusterman MA, Reece GP, Miller MJ, Evans GR, Robb GL, Baldwin BJ. Choice of flap and incidence of free flap success. *Plast Reconstr Surg* 1996;**98**:459-63.
- 7 Robinson DW, MacLeod A. Micro vascular free jejunum transfer. *Br J Plast Surg* 1982;**35**:258-67.
- 8 Fang T, Zhang EW, Lineaweaver WC, Zhang F. Recipient vessels in the free flap reconstruction around the knee. *Ann Plast Surg* 2013;**71**:429-33.
- 9 Gravvanis A, Kyriakopoulos A, Kateros K, Tsoutsos D. Flap reconstruction of the knee: A review of current concepts and a proposed algorithm. *World J Orthop* 2014;**5**:603-13.
- 10 Tsao CK, Chen HC, Chuang CC, Chen HT, Mardini S, Coskunfirat K. Adequate Venous Drainage. *Ann Plast Surg* 2004;**53**:229-234.
- 11 Chia HL, Wong CH, Tan BK, Tan KC, Ong YS. An algorithm for recipient vessel selection in microsurgical head and neck reconstruction. *J Reconstr Microsurg* 2011;**27**:047-056.
- 12 Sarukawa S, Asato H, Okazaki M, Nakatsuka T, Takushima A, Harii K. Clinical evaluation and morbidity of 201 free jejunal transfers for

- oesophagopharyngeal reconstruction during the 20 years of 1984–2003. *Scand J Plast Reconstr Surg Hand Surg* 2006;**40**:148-52.
- 13 Walker R, Praveen P, Parmar S, Martin T, Anand R, Simms M. The jejunal free flap in reconstruction of hypopharyngeal defects – the Birmingham experience. *Br J Oral Maxillofac Surg* 2010;**48**:S15.
 - 14 Reece GP, Schusterman MA, Miller MJ, Kroll SS, Robb GL, Baldwin BJ, Luethcke DR. Morbidity and functional outcome of free jejunal transfer reconstruction for circumferential defects of the pharynx and cervical esophagus. *Plast Reconstr Surg* 1995;**96**:1307-16.
 - 15 Hayashi A, Natori Y, Komoto M, Matsumura T, Horiguchi M, Yoshizawa H, Iwanuma Y, Tsurumaru M, Kajiyama Y, Mizuno H. Microvascular Reconstruction of Free Jejunal Graft in Larynx-preserving Esophagectomy for Cervical Esophageal Carcinoma. *Plast Reconstr Surg Glob Open* 2016;**4**:e632.
 - 16 Ciudad P, Agko M, Manrique OJ, Date S, Kiranantawat K, Chang WL, Nicoli F, Lo Torto F, Maruccia M, Orfaniotis G, Chen HC. The retrograde transverse cervical artery as a recipient vessel for free tissue transfer in complex head and neck reconstruction with a vessel-depleted neck. *Microsurgery* 2017;**37**:902-9.
 - 17 Baj A, Bolzoni A, Torretta S, Pignataro L. Arterial microanastomoses on the reverse flow of the internal carotid artery reverse flow: an extreme solution in free-flap revascularisation. How we do it. *Acta Otorhinolaryngol Ital* 2014;**34**:368-71.
 - 18 Müller DF, Lohmeyer JA, Zimmermann A, Siewert JR, Kovacs L, Machens HG, Biemer E. The carotid artery as recipient vessel: troubleshooting for free jejunal transfer after esophagectomy in preradiated patients. *Chirurg* 2011;**82**:670-4.
 - 19 Kim SH, Kim HK, Kim K, Shim YM. Outcome of free jejunal transfer using the end-to-side arterial anastomosis technique as a pharyngo-oesophageal substitute: a 15-year experience. *Eur J Cardio-Thoracic Surg* 2013;**44**:520-524.
 - 20 Lee H-S, Park SY, Jang H-J, Kim MS, Lee JM, Zo JI. Free jejunal graft for esophageal reconstruction using end-to-side vascular anastomosis and extended pharyngo-jejunoscopy. *Ann Thorac Surg* 2012;**93**:1850-4.
 - 21 Salamoun W, Swartz WM, Johnson JT, Jones NF, Myers EN, Schramm VL Jr, Wagner RL. Free Jejunal Transfer for Reconstruction of the Laryngopharynx. *Otolaryngol Neck Surg* 1987;**96**:149-50.
 - 22 Venkatramani H, Sabapathy SR, Nayak S. Free-flap cover of complex defects around the knee using the descending genicular artery as the recipient pedicle. *J Plast Reconstr Aesthetic Surg* 2014;**67**:93-98.
 - 23 Knobloch K, Herold C, Vogt P. Free latissimus dorsi flap transfer for reconstruction of soft tissue defects of the lower extremity. *Oper Orthop Traumatol* 2012;**24**:122-30.
 - 24 Fisher J, Macht S, Lange T. Latissimus dorsi myocutaneous free flap vascularised by an extra-anatomic graft in the lower limb following traumatic disruption of the popliteal artery. *Br J Plast Surg* 1983;**36**:480-2.
 - 25 Grieb G, Alharbi Z, Simons D, Tsolakidis S, Stromps JP, Piatkowski A, Fuchs P, Pallua N. Abdominal aorta as a recipient artery: Using a free latissimus dorsi myocutaneous flap to close hip and pelvic defects. *Int J Surg Case Rep* 2013;**4**:23-5.